

Monitoring Western Corn Rootworm Activity in Soybeans to Predict Rootworm Injury in First-Year Corn

Ronald B. Hammond, Andy Michel and James B. Eisley, Department of Entomology, The Ohio State University

Western corn rootworm (WCR) (fig. 1), which spread across Ohio during the 1970s from its western origin, has been a problem on field corn in much of Ohio. Historically, injury caused by this species was primarily limited to corn following corn because adult rootworm beetles only deposited their eggs in plantings of corn. The eggs hatched the following spring into larvae that fed on the root systems of corn planted after corn (fig. 2), thus causing the injury that led to yield losses.



Figure 1. Western corn rootworm adult

In the late 1980s, cases of rootworm injury on first-year corn by the WCR were identified in northeastern Illinois and soon thereafter in northwestern Indiana. In this case, adult rootworms were depositing eggs in fields other than corn fields, most often soybean fields. The eggs then hatched the following year in corn planted after soybeans. This new egg-laying behavior enabled populations of WCR to thrive under a system of intense two-year rotations of corn and soybeans. This variant of the WCR spread throughout Illinois and Indiana during the 1990s, especially in an eastward direction towards the Indiana-Ohio border. By 1996, OSU Extension entomologists had confirmed that the new variant of WCR

had become established in Ohio counties bordering Indiana. More recent observations indicate that it has become a more common occurrence in western Ohio, and is now present in the central part of the state.



Figure 2. Root injury

Prediction of WCR Injury in First-Year Corn

The potential risk for rootworm injury in first-year corn can be predicted by monitoring WCR beetle activity in soybeans during the previous season. Adult rootworm beetle activity can be measured by the use of yellow sticky traps (fig. 3). The preferred method for monitoring rootworm beetle activity in soybeans is the use of the Pherocon AM yellow sticky trap.



THE OHIO STATE UNIVERSITY

COLLEGE OF FOOD, AGRICULTURAL,
AND ENVIRONMENTAL SCIENCES

extension.osu.edu

agmr.osu.edu



Figure 3. Yellow sticky trap

When using yellow sticky traps to monitor beetle activity, 6 traps should be stationed in soybeans at least 100 feet into the field and from each other. The traps should be placed in a soybean field by the third week in July, when adult rootworm beetles are beginning to emerge from cornfields. The traps should be replaced at 7-day intervals, with trapping continuing for 6 weeks. Trap monitoring of rootworm beetle activity should be terminated around early September.

The average number of beetles collected per trap per day per sampling period (7 days) over the entire monitoring period provides an index of adult rootworm beetle activity. In general, an average collection of 5 adult beetles per trap per day for any sampling period is an indication of potential economic rootworm injury if corn is planted in the field the following season.

See Ohio State University Extension Bulletin 545, *Control of Insect Pests of Field Crops*, for those insecticides labeled for corn rootworms, or for all insecticides labeled on corn. Bulletin 545 can be accessed at <http://entomology.osu.edu/ag/>.

This publication contains pesticide recommendations that are subject to change at any time. These recommendations are provided only as a guide. It is always the pesticide applicator's responsibility, by law, to read and follow all current label directions for the specific pesticide being used. Due to constantly changing labels and product registration, some of the recommendations given in this writing may no longer be legal by the time you read them. If any information in these recommendations disagrees with the label, the recommendation must be disregarded. No endorsement is intended for products mentioned, nor is criticism meant for products not mentioned. The authors, Ohio State University Extension, and the Ohio Agricultural Research and Development Center assume no liability resulting from the use of these recommendations.

Additional information is available from your local Extension office or The Ohio State University Entomology Agronomic Crops Insects web site (<http://entomology.osu.edu/ag/>).

Ohio State University Extension embraces human diversity and is committed to ensuring that all research and related educational programs are available to clientele on a nondiscriminatory basis without regard to age, ancestry, color, disability, gender identity or expression, genetic information, HIV/AIDS status, military status, national origin, race, religion, sex, sexual orientation, or veteran status. This statement is in accordance with United States Civil Rights Laws and the USDA.

Keith L. Smith, Associate Vice President for Agricultural Administration; Associate Dean, College of Food, Agricultural, and Environmental Sciences; Director, Ohio State University Extension; and Gist Chair in Extension Education and Leadership.

For Deaf and Hard of Hearing, please contact Ohio State University Extension using your preferred communication (e-mail, relay services, or video relay services). Phone 1-800-750-0750 between 8 a.m. and 5 p.m. EST Monday through Friday. Inform the operator to dial 614-292-6181.

Copyright © 2014, The Ohio State University