

Black Cutworm on Corn

Ronald B. Hammond, Andy Michel, and James B. Easley, Department of Entomology, The Ohio State University

Loss of stand prior to emergence or due to above-ground cutting or below-ground tunneling injury often indicates presence of black cutworm (BCW) (fig. 1). Adult moths (fig. 2) of the BCW overwinter in the south and migrate north with the prevailing spring weather fronts. The intensity and distribution of the migratory BCW flight may differ from year to year due to this migration. Some regions such as the Ohio River valley often exhibit higher levels of BCW activity than other regions such as the areas bordering the Great Lakes. Within a region, migrating BCW moths tend to be attracted to fields having a significant ground cover of winter annuals such as chickweed at the time of peak migratory flights. Since the accumulation of winter annuals is often associated with reduced tillage or no-tillage systems, the incidence of BCW infestations tends to be linked to tillage practices and weed management. Late tillage prior to corn planting may not reduce BCW infestations if significant weed cover was present at the time of peak egg laying by migratory BCW moths. Thus, reduced tillage or no-tillage corn may be equally susceptible to BCW infestation if the corn crop follows a no-tillage corn or soybean crop that enabled development of a significant stand of winter annuals, which are attractive sources for egg laying by migratory BCW moths. In terms of weed management, fall tillage can give growers a start in providing weed-free fields in the spring.

Eggs are laid on these weeds in the spring, and developing larvae move to corn when the weeds are killed with tillage or herbicides. BCW larvae pass through seven instars. Early first, second, and third BCW instars are rarely detected in the field, although such early instars may cause slight foliar feeding. Above-ground cutting of corn seedlings (fig. 3) or below-ground tunneling of corn (fig. 4) is generally linked to BCW larvae in the fourth and later instars, which range in length from 1/2 to 2 inches. Stand loss of corn is generally associated with below-ground feeding injury, which occurs below the growing point. Corn plants are most vulnerable to cutworm injury following emergence, and susceptibility to injury declines as the corn plant approaches the 6th leaf stage of development. Plants cut off above the growing point tend to recover.

Symptoms, Assessment, and Scouting

Fields particularly heavy with broadleaf weeds, especially chickweed, should be monitored closely from planting until corn reaches the 6th leaf stage. When infestations of BCW are detected, the proportion of stand exhibiting injury is an indication of additional injury that may be forthcoming. Depending on the stages of cutworm and corn development present, 1 to 3 additional plants may be affected for every plant exhibiting fresh injury up to the 4th leaf stage. Thus, rescue treatment



Figure 1. Black cutworm larva



Figure 2. Black cutworm adult



THE OHIO STATE UNIVERSITY

COLLEGE OF FOOD, AGRICULTURAL,
AND ENVIRONMENTAL SCIENCES

extension.osu.edu

agmr.osu.edu



Figure 3. Above-ground feeding injury



Figure 4. Below-ground feeding injury

should be based on accurate estimates of stand injury and presence of cutworms in the early to mid-larval stages.

Check 20 plants in 5 locations weekly for cutworm injury beginning after corn emergence. Determine percent of plants being injured and collect a number of cutworm larvae to determine the predominant stage of larval development. Rescue treatment is warranted when (1) 3 percent or more of the plants have been cut or tunneled, (2) corn plants are in the 2nd- to 6th-leaf stage, and (3) larvae are still 1 inch or less in length. If a rescue treatment is required, growers have numerous insecticide treatment options.

Management

Because of the relationship between BCW and weed infestations, early tillage and fall and/or spring weed control will reduce the incidence of cutworm infestations. Weed control can be especially useful if done as a fall management program. Conversely, reduced tillage or no-tillage fields with substantial broadleaf weed

infestations may warrant a preventive treatment prior to emergence. Such a treatment might be especially critical if adult cutworm flight is occurring within a region. Preventive control can be achieved by at-planting or pre-emergent use of various insecticides.

Corn hybrids with the Herculex I transgenic trait (including Herculex Xtra) will also control BCW larvae in addition to corn borers. However, hybrids containing YieldGard and Agrisure traits do not offer BCW control. Although the various neonicotinoid seed treatments, Cruiser and Poncho, have BCW on their label, our experience, as well as the experience of others, suggests they do not offer acceptable control. Thus, hybrids treated with these seed treatments that are not from the Herculex I family should be monitored for BCW in the spring.

See Ohio State University Extension Bulletin 545, *Control of Insect Pests of Field Crops*, for those insecticides labeled for black cutworm, or for all insecticides labeled on corn. Bulletin 545 can be accessed at <http://entomology.osu.edu/ag/>.

This publication contains pesticide recommendations that are subject to change at any time. These recommendations are provided only as a guide. It is always the pesticide applicator's responsibility, by law, to read and follow all current label directions for the specific pesticide being used. Due to constantly changing labels and product registration, some of the recommendations given in this writing may no longer be legal by the time you read them. If any information in these recommendations disagrees with the label, the recommendation must be disregarded. No endorsement is intended for products mentioned, nor is criticism meant for products not mentioned. The authors, Ohio State University Extension, and the Ohio Agricultural Research and Development Center assume no liability resulting from the use of these recommendations.

Additional information is available from your local OSU Extension office or The Ohio State University Entomology Agronomic Crops Insects web site (<http://entomology.osu.edu/ag/>).

Ohio State University Extension embraces human diversity and is committed to ensuring that all research and related educational programs are available to clientele on a nondiscriminatory basis without regard to age, ancestry, color, disability, gender identity or expression, genetic information, HIV/AIDS status, military status, national origin, race, religion, sex, sexual orientation, or veteran status. This statement is in accordance with United States Civil Rights Laws and the USDA.

Keith L. Smith, Associate Vice President for Agricultural Administration; Associate Dean, College of Food, Agricultural, and Environmental Sciences; Director, Ohio State University Extension; and Gist Chair in Extension Education and Leadership.

For Deaf and Hard of Hearing, please contact Ohio State University Extension using your preferred communication (e-mail, relay services, or video relay services). Phone 1-800-750-0750 between 8 a.m. and 5 p.m. EST Monday through Friday. Inform the operator to dial 614-292-6181.

Copyright © 2014, The Ohio State University